

Case Study II: Models of transport, waves, and shallow waters

Review questions and exercises

1 Modeling

1.1 Review questions

1. Specify the conservative form, the primitive form, and the integral form of a scalar conservation law in one space dimension.
2. What is the significance of the characteristics associated with a conservation law?
3. Sketch the characteristics for Burgers equation associated with the initial condition u_0 in figure 1, and sketch qualitatively the initial development of the solution with this boundary condition. Note the different behaviors to the right and left of $x = 0$.
4. Under which condition is the momentum conserved in a fixed amount of fluid?

1.2 Exercises

1. Define the characteristics of the conservation law $u_t + f(u)_x = 0$.
2. Let $u(x, t)$ be a velocity field computed by solving a scalar conservation law in one space dimension; $u > 0$ corresponds to a velocity directed along increasing x coordinate. Write up the ODE satisfied by the x -coordinate $p(t)$ of a massless particle that starts at position x_0 and is transported by the velocity field along the x axis.
3. Rewrite following equations in conservative form and specify the flux function:

(a)

$$u_t + uu_x = 0 \tag{1}$$

(b)

$$u_t + uu_x + vu_y = 0, \tag{2}$$

under the condition that $u_x + v_y = 0$.

(c)

$$uu_t + u^2 u_x = 0, \tag{3}$$

as a conservation law in u^2 .

Remark 1. Note that equation (3) is obtained by multiplying equation (1) by u . Equations (1) and (3) are thus equivalent for smooth solutions. However, their conservative forms will be different and corresponding shock propagation speeds will differ!

4. For arbitrary twice differentiable function f and g , show that

$$p(x, t) = f(x + ct) + g(x - ct) \tag{4}$$

is a solution to the wave equation

$$p_{tt} - c^2 p_{xx} = 0. \tag{5}$$

From expression (4), conclude the form of the characteristics associated with the wave equation (5).

5. (a) Rewrite the wave equation (5) as a system of conservation laws

$$\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = \mathbf{0} \tag{6}$$

in $\mathbf{u} = (u, v)$, where $u = -cp_x$ and $v = p_t$, and specify the 2-by-2 matrix \mathbf{A} in the representation $\mathbf{f}(\mathbf{u}) = \mathbf{A}\mathbf{u}$ of the flux function.




FIGURE 1. An initial condition

- (b) By defining variables w and z as linear combinations of the above u and v , rewrite equation (6) as two uncoupled scalar transport equations, representing transport in the positive and negative x -direction respectively. (These are called *characteristic variables*).
6. Consider Burgers equation with smooth initial data $u_0(x)$,

$$\begin{aligned} u_t + \frac{1}{2}(u^2)_x &= 0 & t > 0, \\ u(x, 0) &= u_0(x). \end{aligned} \quad (7)$$

Claim: The characteristics will cross (and thus a shock form) if $u'(x) < 0$ at any point x , and the time for the first occurrence of crossing characteristics is

$$T^* = -\frac{1}{\min_x u'(x)}. \quad (8)$$

- (a) Compute T^* for the initial condition (figure 1)

$$u_0(x) = \begin{cases} 1 + \cos \pi x & -1 \leq x \leq 1, \\ 0 & \text{otherwise.} \end{cases} \quad (9)$$

- (b) Prove the above claim.

7. Generalize to two space dimensions x, y , the derivation of the shallow water equations over a planar bottom at $z = 0$. In this case, the unknown functions will be the water level $h(x, y)$ and components $u(x, y), v(x, y)$ of the horizontal velocity vector $\mathbf{u} = (u, v, 0)$. *Hint:* There will be two moment balance laws, one for ρu and one for ρv .

2 FVM

2.1 Review questions

1. The family of conservative, explicit finite-volume schemes associated with conservation law $u_t + f(u)_x = 0$ that we consider is

$$u_i^{n+1} = u_i^n - \frac{\Delta t}{\Delta x} (F_{i+1/2}^n - F_{i-1/2}^n) \quad (10)$$

- (a) What is approximated by the quantities u_i^n ?
- (b) What is the numerical flux function $F_{i+1/2}$ for the upwind scheme when $f(u) = cu$ in the case (i) $c > 0$, and (ii) $c < 0$?
2. Give (a least) one good and one bad property with the upwind scheme, the Lax–Friedrich method, and the Richtmyer two-step Lax–Wendroff method.
3. What is meant by the CFL condition for a numerical scheme associated with a conservation law?
4. Are we guaranteed that a numerical scheme is stable if it satisfies the CFL condition? Can a numerical scheme violate the CFL condition and still be stable?

2.2 Exercises

1. Show that the upwind numerical flux for the particular case $f(u) = cu$ can be written

$$F_{i+1/2}^n = \frac{1}{2}(f(u_{i+1}^n) + f(u_i^n)) - \frac{|c|}{2}(u_{i+1}^n - u_i^n), \quad (11)$$

and that the update formula (10) with numerical flux (11) becomes

$$u_i^{n+1} = u_i^n - \frac{\Delta t}{2\Delta x} [f(u_{i+1}^n) - f(u_{i-1}^n)] - \frac{|c|\Delta t}{2\Delta x} [2u_i^n - u_{i-1}^n - u_{i+1}^n]. \quad (12)$$

Remark 2. Expression $(f(u_{i+1}) - f(u_{i-1})) / (2\Delta x)$ constitutes a *central difference approximation* of $f(u)_x$, and $-(2u_i - u_{i-1} - u_{i+1})\Delta x^2$ a finite-difference approximation of $u''(x_i)$. Expression (12) can thus be viewed as a discretization of the equation

$$u_t + f'(u)_x = |c|\Delta x u''. \quad (13)$$

with a central difference approximation of $f(u)_x$. Hence, an *upwind* discretization of $u_t = f(u)_x$ for $f(u) = cu$ is equivalent to a *central* discretization of the conservation law with an added *artificial dissipation* (the last term in expression (12) and the right side of equation (13)). This interpretation is consistent with the very diffusive properties of the upwind discretization.

2. A common implementation of the upwind scheme for nonlinear f is to use formula (11) and substitute c with an approximation of f' at the interface. Verify that we obtain the upwind scheme for Burgers equation when using the numerical flux (11) with $c = (u_i + u_{i+1})/2$.
3. Show that the Lax–Friedrich method for the particular case $f(u) = cu$ will be the same as the upwind method for a particular choice of time step Δt .
4. Show that the upwind scheme exactly solves the transport equation $u_t + cu_x = 0$ if the time step is chosen at the stability limit for the CFL condition. (This property does *not* hold for nonlinear conservation laws!)